



International Conference Fund
Funded by the NZ Government.
Administered by the Royal Society of NZ

UV radiation and its effects

– an update 2010

Report of the NIWA UV Workshop
Queenstown, 7–9 May 2010

Organising Committee

Dr Richard McKenzie, NIWA, Lauder (Convenor)
Assoc. Prof. Robert Scragg, University of Auckland (Co-convenor)
Dr Judith Galtry, Cancer Society (Co-convenor)
Graeme Strang, NIWA, Lauder (Registrations)

Sponsors

National Institute of Water & Atmospheric Research
Cancer Society of New Zealand
The Royal Society of New Zealand
Ministry of Health
Health Sponsorship Council/Sunsmart
Foundation for Research Science & Technology

NIWA Information Series No.77
2010

**Published by
National Institute of Water & Atmospheric Research
(NIWA)
Wellington**

**© NIWA
2010**

UV radiation and its effects – an update 2010

**ISBN 978-0-478-23289-9
ISSN 1174-264X**

The previous workshop in this series was published as
UV radiation and its effects – an update 2006
RSNZ Miscellaneous Series 68
ISBN 1-877264-20-2
ISSN 0111-3895

Contents

UV Workshop Overview	Richard McKenzie	5
Ozone: Past, Present and Future	Greg Bodeker	7
The Antarctic Ozone Hole: Didn't we fix that already?	Stephen Wood	9
Developing UV monitoring with CCD technology	Bruce Forgan and Peter McGlynn	11
Spectral measurements of solar UV at several altitudes under Australian conditions	Stuart Henderson et al.	13
Effect of snow albedo and topography on UV radiation	Julian Gröbner and Gregor Hülsen	15
Environmental impacts of ozone depletion and its interactions with climate change	Janet Bornman	17
Proxy methods to estimate UV	Ken Ryan and Allan Burne	19
UV radiation in New Zealand: implications for grape quality	R Hofmann et al.	21
Improved UVI products at NIWA	Richard Turner	23
UV dosimeter based on polyphenylene oxide for the measurement of UV exposures to plants and humans over extended periods	Alfio Parisi et al.	25
Electronic UV dosimeters for research and education	Martin Allen	27
Personal exposures to UV radiation in New Zealand	Ben Liley et al.	29
Personal UV dosimeter badges: Mark II	David Sherman	31
Calibration of solaria	Paul Johnston and Richard McKenzie	33
Erythema versus vitamin D production from sunlight and solaria	Richard McKenzie et al.	35
The introduction of compact fluorescent lights (CFLs) and the impact of UVR emissions on photosensitive people	John Javorniczky et al.	37
Solaria in Australia and the standards making process	Peter Gies et al.	39
Sun bed policy: an international perspective	Craig Sinclair	41
Photoageing – what is it all about?	Marius Rademaker	43
The role of built environment in modifying UV exposure	Christina Mackay	45
Correlates of sun exposure in people aged 45 and older in NSW	Anne Kricker et al.	47
Vitamin D, past sun exposure & skin phenotype in risk of central nervous system demyelination	Robyn Lucas	49
Vitamin D3, the skin immune system and cutaneous carcinogenesis	Roslyn Malley et al.	51
Considerations for increasing vitamin D in the food supply	Julia Edmonds	53
Vitamin D and cardiovascular disease: are we at a tipping point?	Robert Scragg	55
Factors associated with body coverage and personal UV radiation dose: preliminary findings	Anthony Reeder et al.	57
Vitamin D and its association with personal UV exposure	Alistair Stewart and Robert Scragg	59
Vitamin D in Brisbane office workers	Lan Vu et al.	61
Action spectra for UV effects – UV filtering by human skin	Merve Meinhardt-Wollweber	63
Seasonal vitamin D	Michael Kimlin	65
The increasing incidence of melanoma: the role of MelNet	Graham Stevens	67
Maximising protection — sun protective fabric and clothing	Cheryl Wilson	69
The Ultraviolet Index: health promotion tool or “poisoned chalice”?	Judith Galtry	71

Developing an effective UV Alert: a qualitative study	Jennifer Makin et al.	73
Communicating UVR information to the public	Rebecca Gray and Wayde Beckman	75
Burden of disease associated with low vitamin D status in New Zealand	Robert Scragg et al.	77
The estimated costs – economic and human – of skin cancers to New Zealand	Des O’Dea	79
 Posters		
Comparison between UV irradiances in Christchurch city and the Port Hills	Richard McKenzie et al.	81
Two decades of spectral UV measure	Michael Kotkamp et al.	82
Characterisation of AlGaIn detectors for UV measurements	Neil Swift et al.	84
Processing and results from RB meter networks in New Zealand	John Sansom	86
Analysis of dosimeter badge data	James Liley and Ben Liley	88
Dermatological lamps as a source of vitamin D	Roslynd Winters	90
Understanding UV reflection in the urban environment	Joanna Turner and Alfio Parisi	92
Neonatal exposure to solar simulated ultraviolet radiation leads to deviation of immune development	Heather McGee et al.	94
Estimation of the benefits of raising mean serum 25-hydroxyvitamin D levels to 100-115 nmol/L world wide	William Grant	96
High-risk patients self-select for melanoma screening	Amanda Oakley and Marius Rademaker	98
Vitamin D status of a Dunedin sample, 2008–09: preliminary results	Janet Jopson et al.	100
Sun protection and vitamin D	Louise Reiche	102
UV Index variations (1960–2006) and melanoma skin cancer incidence in Australia	Lemus-Deschamps and Jen Makin	104
Action spectrum for vitamin D synthesis	William Olds et al.	106
Effects of clouds and aerosols on UV radiation measured at Lauder, New Zealand	Jordi Badosa et al.	108
Determinants of vitamin D in a multi-ethnic sample of Auckland residents	Sofia Nessvi et al.	110
Thinner melanomas detected by digital monitoring	Marius Rademaker and Amanda Oakley	112
Dermoscopic global pattern criteria in practice	Marius Rademaker and Amanda Oakley	114
Inter-observer variability of teledermoscopy	Marius Rademaker et al.	116
Seasonal variation in vitamin D levels and UVB exposure in Christchurch	C. Florkowski et al.	118
 Participants		 120

Action spectra for UV effects – UV filtering by human skin

Merve Meinhardt-Wollweber

Leibniz Universität Hannover, Institute of Biophysics, Hannover, Germany

Abstract. Human skin is the interface between man and his environment. One of the environmental factors it has to deal with is ultraviolet radiation. Ultraviolet radiation may cause negative effects such as erythema, skin ageing, skin cancer, etc. Not surprisingly, the epidermis – the upper layer of the skin – acts as a natural UV filter and may thicken or develop a stronger pigmentation as an adaptation to excess UV exposure. On the other hand, UV radiation induces photoproduction of previtamin D₃ in the skin and thus provides the major natural source for vitamin D₃ for humans. Thus, it seems reasonable that skin has not evolved as a mere sun block but as a sun screen which is even adaptable to a certain extent. Using optoacoustics, we have determined the wavelength dependent absorption coefficients of human skin in the UVB and UVA-II. Based on these data for the UV filtering properties of human skin, we calculated the biological effectiveness of solar radiation at different depths within the epidermis and found that UV adaptation more effectively shields the skin from erythemally active radiation than from vitamin D producing radiation. Thus, adaptation to high UV exposure tends to choose beneficial over detrimental UV effects.

Optoacoustic study

Optoacoustics is a fairly new method for studying the optical properties of human skin *in vivo*. Pulsed laser radiation is used to illuminate a sample of the skin. The radiation is distributed in the sample according to its optical properties and absorbed radiation energy is converted to pressure. This pressure profile (high pressure where a lot of radiation has been absorbed and low pressure at volumes with low absorption) is released as an ultrasound pulse and can be measured above the skin. The initial light distribution in the sample and its optical properties can be deduced from this ultrasound pulse.

We measured the absorption spectra of human skin in a study in 20 subjects belonging to skin phototypes I-IV (phototype I: n=3, phototype II: n=6, phototype III: n=6, phototype IV: n=5). Absorption coefficients were determined in the range from 290 nm to 341 nm (3 nm steps) on the inner and outer side of the forearm and the ball of the thumb. The study was conducted in the summertime and 12 subjects reported sunseeking behaviour whereas the other 8 subjects said they were office workers and tended to avoid sun exposure.

The optical properties that human skin develops depend on individual native and environmental parameters. Environmental effects may be facultative pigmentation and a thickened horny layer in response to and as an adaptation to higher UV exposure. Because the skin on the inner side of the forearm is relatively thin and normally only weakly exposed to daily sunlight, it is expected to show nearly the native properties of individual skin. On the outer side of the forearm, the skin is much more

strongly exposed to sunlight and UV radiation but the skin structure is still similar to that on the volar side. Comparative analysis of these two sites allows insight into how UV adaptation may affect the optical properties of the skin.

Optical properties of skin

Figure 1 shows the absorption spectra of human skin at different skin sites: inner and outer side of the forearm – i.e., skin with low and high natural UV exposure – and the ball of the thumb where the skin's horny layer is especially thick. Data for the outside of the forearm is shown as mean as well as split into the high and low UV exposure group. Remarkably, the horny layer is a potent UVB screen but is fairly transparent to UVA radiation. Adaptation to higher UV exposure results in higher absorption coefficients, especially in the UVB range, probably due to epidermal thickening and pigmentation.




Figure 1. UV absorption spectra from different sites of human skin: inner side of the forearm (black circles), outer side of the forearm (mean: bi-colour diamonds, low UV exposure group: downward triangles, high UV exposure group: upward triangles) and ball of the thumb/horny layer (squares).




Figure 2. Penetration depths (1/e-level) of UV radiation for human skin at the volar side of the forearm (squares), dorsal side of the forearm (circles), and at the ball of the thumb (triangles).

Figure 2 shows what these absorption spectra mean in terms of penetration depth. In normal skin, as on the forearm, UV radiation is strongly attenuated. Only 40% or less of the incoming radiation reaches a depth of 60 µm – halfway through the viable epidermis. The horny layer very effectively reduces the penetration depth for UVB radiation. A thickened horny layer thus provides an effective screen for the living cells underneath it. On the other hand, it is ineffective in blocking UV radiation.

UV filtering and UV effects

To assess the UV filtering potential of human skin, we calculated how much of the diffuse solar spectral irradiance weighted by the action spectrum for erythema or previtamin D₃ formation penetrates 20 and 100 µm deep into human skin based on its optoacoustically measured optical properties.

Applying a simple model like the Lambert-Beer law, the global spectral irradiance $E(\lambda)$ is attenuated exponentially with increasing depth d according to the skin's optical properties (μ_a). This attenuated spectrum is weighted by the respective action spectrum $A(\lambda)$ and integrated over the relevant wavelength range yielding the biologically effective irradiance at the respective depth:

$$E_{bio}^{skin} = \int_{290nm}^{345nm} E(\lambda)e^{-\mu d} * A(\lambda)d\lambda$$

Figure 3 shows the difference between $E_{erythema}$ ($isE_{erythema}$) and $E_{previtaminD3}$ ($isE_{previtaminD3}$) for two depths within the skin, weakly pigmented vs. UV adapted skin, and for global spectral irradiances from SZA 30°, 50° and 70° as would be expected for summer, autumn/spring, and winter at temperate latitudes. Only when the sun is low, vitamin D₃ is produced throughout the skin before the minimal erythemal dose is reached. Erythema outweighs vitamin D₃ production for the rest of the year in weakly pigmented skin. The cumulative UV-screening effect of the skin chromophores becomes, of course, more and more effective with increasing path length – i.e., depth – in the skin. At the basal layer, where protection from detrimental UV effects is most important, the full impact of the filtering properties of UV adapted skin is unveiled:

100 µm of UV adapted skin can shift the dominant effect of UV radiation from erythema to vitamin D₃ production.




Figure 3. Difference between the biologically effective irradiance of one detrimental UV effect ($E_{erythema}$ ($isE_{erythema}$)) and potentially beneficial UV-induced vitamin D₃ synthesis ($E_{previtaminD3}$ ($isE_{previtaminD3}$)). Combinations of solar spectral irradiance from three different SZA and two UV adaptation status of the skin are compared. The upper bar in each row is for a depth of 20 µm within the skin and the lower bar (in red) shows the situation at a depth of 100 µm near the basal layer at the dermal-epidermal-junction.

It is important to note, however, that the action spectra used for this calculation can give only a rough idea of the true in vivo situation. The action spectrum for previtamin D₃ production for example marks the dose where 5% of provitamin D₃ is converted to previtamin D₃. It is not clear what this means in terms of vitamin D₃ sufficiency. Besides, all action spectra are based on single dose administration, but the factor of time is very important in the in vivo situation. For example, repair mechanisms may take their time to counteract detrimental UV effects. On the other hand vitamin D₃ may just accumulate over time.

References

- Anders, A., Altheide, H.-J., and Tronnier, H., 2002. Action Spectroscopy of Human Skin with Tunable Lasers, CIE (research note), 148, 1-9.
- Bernhard, G., Mayer, B., and Seckmeyer, G., 1997. Measurements of spectral solar UV irradiance in tropical Australia, J. Geophys. Res., 102(D7), 8719-8730.
- Bouillon, R., Eisman, J., Garabedian, M., Holick, M., Kleinschmidt, J., Suda, T., Terenetskaya, I., and Webb, A., 2006. Action spectrum for the production of previtamin D3 in human skin, CIE (technical report), 174, 1-12
- Meinhardt, M., Krebs, R., Anders, A., Heinrich, U., and Tronnier, H., 2008. Effect of ultraviolet adaptation on the ultraviolet absorption spectra of human skin in vivo. Photoderm. Photoimmunol. Photomed., 24, 76-82.
- Meinhardt, M., Krebs, R., Anders, A., Heinrich, U., and Tronnier, H., 2008. Wavelength dependent penetration depths of ultraviolet radiation in human skin. J. Biomed. Opt. 13(4), 044030.
- Meinhardt, M., Krebs, R., Anders, A., Heinrich, U., and Tronnier, H., 2009. Absorption spectra of human skin in vivo in the ultraviolet wavelength range measured by optoacoustics. Photochem. Photobiol., 85, 70-77.
- Meinhardt-Wollweber, M., Krebs, R., Anders, A., Heinrich, U., and Tronnier, H., 2009. Optoacoustics – Application in Photodermatology. IFSCC magazine 12(4).